

2017 AUTUMN VARIETY RATINGS FACT SHEET

BLACKLEG MANAGEMENT GUIDE



Quantify the risk, paddock by paddock

Blackleg can cause severe yield loss, but can be successfully managed. Use this guide to determine whether you are in a high-risk situation and what practices you can change to reduce or prevent yield loss from blackleg. Follow the four steps, in sequence, below.

KEY POINTS

- **Never sow your canola crop into last year's canola stubble.**
- **Monitor your crops in Spring to determine yield losses in the current crop.**
- **Choose a cultivar with adequate blackleg resistance for your region.**
- **Relying only on fungicides to control blackleg poses a high risk of fungicide resistance.**
- **If your monitoring has identified yield loss and you have grown the same cultivar for three years or more, choose a cultivar from a different resistance group.**

Blackleg is a sexually reproducing pathogen that will overcome cultivar resistance genes. Fungal spores are released from canola stubble and spread extensively via wind and rain splash. The disease is more severe in areas of intensive canola production.

STEP 1: Use Table 1 to determine your farm's blackleg risk

TABLE 1 Regional blackleg factors									
Environmental factors that determine risk of severe blackleg infection	Blackleg severity risk factor								
	High risk			Medium risk			Low risk		
Regional canola intensity (% area sown to canola)	above 20	16–20	15	11–14	11–14	10	6–9	5	below 5
Annual rainfall (mm)	above 600	551–600	501–550	451–500	401–450	351–400	301–350	251–300	below 250
Total rainfall received March–May prior to sowing (mm)	above 100	above 100	above 100	above 100	91–100	81–90	71–80	61–70	below 60

Combined high canola intensity and adequate rainfall increase the probability of severe blackleg infection.

STEP 2: Determine each crop's blackleg severity in Spring

- Assess the level of disease in your current crop. Sample the crop anytime from the end of flowering to windrowing (swathing). Pull 60 randomly chosen stems out of the ground, cut off the roots with a pair of secateurs and, using the reference photos in Table 2 below, estimate the amount of disease in the stem cross-section. Yield loss commonly occurs when more than 50% of the cross-section of the cut stem is discoloured.
- A dark-coloured stem is a symptom of blackleg (Table 2). Stem cankers are clearly visible at the crown of the plant. Severe cankers may cause the plant to fall over as the roots become separated from the stem.
- If you have identified that you are in a high-risk situation (Steps 1 and 2), use Steps 3 and 4 to reduce your risk of blackleg for future seasons.
- If you are in a low risk situation and you have not identified yield loss due to blackleg infection when assessing your crop, continue with your current management practices.

PHOTO: STEVE MARCROFT




TABLE 2 Crop blackleg severity							
High risk		Medium risk			Low risk		
Cankered	100%	80%	60%	40%	20%	0%	

Yield loss occurs when more than half of the cross-section is discoloured.

STEP 3: Management practices can reduce the risk of blackleg infection

If your crop monitoring (see Step 2) showed yield loss (cut stems were >50% discoloured) in the previous year, the following practices can be used to reduce blackleg severity. Complete the following process for each canola paddock to be sown.

- For each of the seven management factors listed below, circle where each canola paddock fits to determine the risk of blackleg. For example, **Blackleg rating**: if your cultivar is ATR-Stingray, circle MR indicating a low risk of blackleg; or **Distance from last year's canola stubble**: if your proposed canola crop is 200 metres away, high risk is indicated.
- Complete all seven management factors to determine which practices are causing increased risk and how they can be reduced. For example, for Distance from last year's canola stubble, choose a different paddock, at least 500m away from last year's stubble, reducing the risk from high to low.

WARNING: 'CANOLA ON CANOLA' WILL CAUSE A SIGNIFICANT YIELD LOSS AND WILL REDUCE THE EFFECTIVE LIFE OF CANOLA CULTIVARS AND FUNGICIDES.

Blackleg management practices that determine risk of blackleg infection, from highest to lowest effectiveness are:

A. BLACKLEG RATINGS

The cultivar blackleg rating is the most important blackleg management tool. If your previous crop had a high level of disease, choose a cultivar with a higher blackleg rating. The 2017 Blackleg Ratings are listed in Table 3 on page 4.

High risk				Medium risk			Low risk	
VS	S-VS	S	MS-S	MS	MR-MS	MR	R-MR	R

*VS = very susceptible; S = susceptible; MS = moderately susceptible; MR = moderately resistant; R = resistant

B. DISTANCE FROM LAST YEAR'S CANOLA STUBBLE

The distance of your current crop to last year's canola stubble will determine disease severity.

NEVER sow your canola crop into last year's canola stubble. Distances from last year's stubble up to 500m will reduce blackleg severity.

High risk				Medium risk			Low risk
0m	100m	200m	300m	400m	500m	>500m	

C. FUNGICIDE USE

Fungicides will provide an economic return only if your crop is at high risk of yield loss. Fungicides complement other management practices, never rely solely on fungicides.

RELIANCE ON FUNGICIDES TO CONTROL DISEASE POSES A HIGH RISK OF FUNGICIDE RESISTANCE.

High risk		Medium risk				Low risk
No fungicide	Foliar applied fungicide	Seed dressing fungicide	Fertiliser applied fungicide	Seed dressing + fertiliser applied fungicide	Seed dressing or fertiliser applied + foliar fungicide	

D. YEARS OF SAME CULTIVAR GROWN

The blackleg pathogen will overcome cultivar resistance genes if the same genes are used each year. By sowing a cultivar based on different resistance genes, the ability of the pathogen to overcome resistance will be reduced. All cultivars have been placed into different blackleg resistance groups based on their resistance complement (see Table 3).

If you have:

- high or increasing levels of blackleg in your crop (from monitoring disease levels each year);
- used the management practices outlined here in Step 3; and
- sown cultivars from the same resistance group in close proximity (within 2km) for three or more years, then sow a cultivar from a different resistance group (see page 4 – Blackleg Resistance Groups).

High risk		Medium risk			Low risk
Sown the same cultivar-resistance group for more than 3 years	Sown the same cultivar-resistance group for 3 years		Sown the same cultivar-resistance group for 2 years	Sown the same cultivar-resistance group the previous year	Sown cultivar from a different resistance group

E. DISTANCE FROM TWO-YEAR-OLD CANOLA STUBBLE

Stubble older than two years produces fewer blackleg spores and will normally have minimal effects on blackleg severity, even where canola is sown into two-year-old stubble. However, two-year-old stubble may cause disease if inter-row sowing canola (see Point F, Canola stubble conservation) or if the cultivar resistance has been overcome.

High risk				Medium risk		Low risk
				0m	100m	250m
						500m
						>500m

STEP 3 CONTINUED PAGE 3

STEP 3 FROM PAGE 2

F. CANOLA STUBBLE CONSERVATION

Stubble destruction is not effective in reducing blackleg infection. Inter-row sowing canola into two-year-old canola stubble where germinating seedlings are immediately next to standing stubble may result in higher levels of blackleg infection.

High risk	Medium risk	Low risk
	Inter-row sowing	Disc tillage Knife point tillage Burning/ burying tillage

G. MONTH SOWN

Canola is most vulnerable to blackleg as a seedling. If crops are sown early into warmer conditions and get through the seedling growth stage quickly, they may escape high blackleg severity.

High risk	Medium risk	Low risk
	June to August	May 15 to 31 May 1 to 14 April 15 to 30

H. DUAL-PURPOSE GRAZING CANOLA

Grazing canola can increase the severity of blackleg in the crop. To minimise any associated reduction in grain yield select a cultivar with a high level of blackleg resistance ($\geq R-MR$), and if using a cultivar with a lower level of resistance, consider use of a fungicide (keeping in mind chemical with-holding periods).

High risk	Medium risk	Low risk
	Grazing canola	

CLICK ON THE FOLLOWING LINK TO ACCESS A WORKSHEET THAT ALLOWS YOU TO VISUALISE THE BLACKLEG RISK OF YOUR INDIVIDUAL Paddock AND HOW YOU CAN REDUCE THIS RISK.

<http://www.grdc.com.au/GRDC-FS-BlacklegManagementGuide>



STEP 4: Blackleg resistance groups

Canola cultivars have different combinations of blackleg resistance genes. Over time, growing cultivars with the same blackleg resistance genes has led to changes in the blackleg pathogen's virulence which has enabled it to overcome cultivar blackleg resistance. By rotating between cultivars with different resistance genes, you can reduce the probability of resistance breakdown and reduce disease severity.

Based on Steps 1 to 3, have you observed increasing blackleg severity and have been growing the same cultivar in close proximity for three years or more?

- **No** - Your current management practices should be sufficient to adequately manage blackleg resistance.
- **Yes** - You may be at risk of the blackleg fungus overcoming the blackleg resistance of your cultivar and it is recommended you grow a cultivar with a different combination of blackleg resistance genes.
- To facilitate this process, all cultivars have been placed into groups (A to H) based on their resistance genes in **Table 3**.
- **You do not need to change resistance groups (cultivars) every year.**

How to use Table 3

1. Identify the resistance group of your previously grown cultivar using the column labelled **Section A – 'Resistance group of cultivar'** (shaded in Light Purple). Note; Some cultivars belong to multiple groups. Some cultivars have not yet been classified and rotation recommendations cannot be made for these cultivars.

Examples: AV-Garnet belongs to Resistance Group A

Hyola® 575CL belongs to resistance Groups B and F

If your previously grown cultivar is not included in Table 3 (page 4), as it is no longer commercially available, refer to Table 4 (page 5).

2. Using **Section B** look down the column with the resistance group of the variety grown previously (e.g. Column A if AV-Garnet was grown previously, or Column BF if Hyola® 575CL was grown previously) to identify cultivars with reduced risk.

■ Green = best possible rotation (no resistance genes in common),

■ Blue = OK rotation (at least 1 resistance gene not in common),

■ Orange = not advised (all resistance genes in common).

Examples: AV-Garnet (Resistance Group A) for 2017 planting – sown after cells shaded ■ Orange (e.g. Pioneer® 45Y88CL) is not recommended, following with anything shaded ■ Blue (e.g. IH30 RR) is okay and anything shaded ■ Green (e.g. ATR-Stingray) is best. Hyola® 575CL (resistance Groups BF) for 2017 planting – sown after cells shaded ■ Orange (e.g. Nuseed Diamond) is not recommended, following anything shaded ■ Blue (e.g. Pioneer 45Y91 CL) is okay and anything shaded ■ Green (e.g. ATR-Mako) is best.

SEE PAGE 4, FOR TABLE 3

TABLE 3 2017 Autumn Blackleg Ratings and Resistance Groups. See page 3, (Step 4) for information on how to use this table.

Variety	2017 Blackleg Rating Bare	2017 Blackleg Rating Jockey	Type	Section A - Resistance group of cultivar	Section B - Resistance group of previous year's cultivar (stubble)											
					A	B	C	AB	AC	ABD	ABF	ABS	ABDF	BF	BC	H
CONVENTIONAL VARIETIES																
SF Brazzil	R-MR		Winter, open pollinated	BC												
Victory® V3002	MR	R	High stability oil, Hybrid	ABF												
Nuseed Diamond [®]	MR		Hybrid	ABF												
AV-Garnet [®]	MS		Open pollinated	A												
TRIAZINE TOLERANT VARIETIES																
Hyola® 350TT	R		Hybrid	ABDF												
Hyola® 650TT	R		Hybrid	ABD												
Pioneer® 44T02 TT [®]	R-MR	R	Hybrid	ABD												
Hyola® 559TT	R-MR		Hybrid	ABD												
Monola® 416 TT	MR		High stability oil, open pollinated	B												
Monola® 515 TT	MR		High stability oil, open pollinated	Different blackleg resistance pattern, effective rotation with existing groups currently unknown												
SF Ignite TT	MR	R	Hybrid	BF												
DG 560TT	MR	R	Hybrid	BF												
DG 670TT	MR	R	Hybrid	BF												
ATR-Mako [®]	MR		Open pollinated	A												
ATR-Stingray [®]	MR		Open pollinated	C												
InVigor T 4510	MR-MS	R	Hybrid	BF												
SF Turbine TT	MR-MS	R-MR	Hybrid	BF												
Pioneer® 45T01 TT [®]	MS	R-MR	Hybrid	AB												
ATR-Wahoo [®]	MS		Open pollinated	A												
ATR-Bonito [®]	MS		Open pollinated	A												
CLEARFIELD SYSTEM VARIETIES																
Hyola® 970CL	R		Winter, Hybrid	H												
Hyola® 575CL	R		Hybrid	BF												
SF Edimax CL	R-MR		Winter, Hybrid	C												
Pioneer® 44Y90 CL	R-MR	R	Hybrid	B												
Pioneer® 44Y89 CL	MR	R	Hybrid	BC												
Victory® V7001 CL	MR	R	High stability oil, Hybrid	ABF												
Pioneer® 45Y91 CL	MR	R	Hybrid	B												
Banker CL	MR-MS		Hybrid	A												
Pioneer® 45Y88 CL	MS	R-MR	Hybrid	A												
Archer	MS		Hybrid	C												
Carbine	MS	MR	Hybrid	A												
ROUNDUP READY VARIETIES																
Nuseed GT53	R		Hybrid	ABDF												
Hyola® 506RR	R		Hybrid	ABD												
Nuseed GT42	R		Hybrid	ABDF												
Hyola® 404RR	R-MR		Hybrid	ABD												
Nuseed GT41	R-MR		Hybrid	ABF												
Pioneer® 44Y24 RR	R-MR	R	Hybrid	C												
Monola® G11	R-MR		High stability oil, Hybrid	ABS												
Hyola® 600RR	MR		Hybrid	ABD												
Pioneer® 43Y23 RR	MR	R	Hybrid	B												
Victory® V5003 RR	MR	R	High stability oil, Hybrid	A												
Pioneer® 45Y25 RR	MR	R	Hybrid	BC												
DG 460RR	MR	R	Hybrid	A												
IH52 RR	MR		Hybrid	AB												

TABLE 3 2017 Autumn Blackleg Ratings and Resistance Groups. See page 3, (Step 4) for information on how to use this table (continued).

Variety	2017 Blackleg Rating Bare	2017 Blackleg Rating Jockey	Type	Section A - Resistance group of cultivar	Section B - Resistance group of previous year's cultivar (stubble)											
					A	B	C	AB	AC	ABD	ABF	ABS	ABDF	BF	BC	H
ROUNDUP READY VARIETIES																
InVigor R 5520P	MR	R	Hybrid	AC												
IH30 RR	MR-MS	R	Hybrid	AB												
IH51 RR	MR-MS	R	Hybrid	A												
ROUNDUP READY and TRIAZINE TOLERANT VARIETIES																
Hyola® 525RT®	MR		Hybrid	ABD												
Hyola® 725RT®	MR		Hybrid	ABD												
Bayer 3000 TR	MS-S	MR	Hybrid	B												

TABLE 4 Resistance groups of cultivars that are no longer commercially available.

CULTIVAR	RESISTANCE GROUP
ATR-Gem	A
Hyola® 50	AD
Hyola® 450TT	ABD
Hyola® 474CL	BF
Hyola® 577CL	B
Monola® 314 TT	Unknown
Monola® 513 GT	A
Nuseed GT50	ABF
Pioneer® Sturt TT	C

BLACKLEG RESISTANCE GROUP MONITORING

Representative cultivars from all blackleg resistance groups are sown in trial sites in all canola-producing regions across Australia and monitored for blackleg severity. These data provide regional information on the effectiveness of each blackleg resistance group and are available on the NVT Online website (www.nvtonline.com.au).

USEFUL RESOURCES

Canola best practice management guide for south-eastern Australia

Availabe Ground Cover Direct, 1800 110 044, www.grdc.com.au/bookshop

Diseases of Canola and their Management: The Back Pocket Guide

www.grdc.com.au/GRDC-BPG-CanolaDiseases

Availabe Ground Cover Direct, 1800 110 044, www.grdc.com.au/bookshop

Australian Oilseeds Federation – Agronomy Centre www.australianoilseeds.com
Ground Cover Supplement (issue 116): Foliar fungal diseases of Pulses and oilseeds

DISCLAIMER

Any recommendations, suggestions or opinions contained in this publication do not necessarily represent the policy or views of the Grains Research and Development Corporation. No person should act on the basis of the contents of this publication without first obtaining specific, independent, professional advice. The Corporation and contributors to this Fact Sheet may identify products by proprietary or trade names to help readers identify particular types of products. We do not endorse or recommend the products of any manufacturer referred to. Other products may perform as well as or better than those specifically referred to. The GRDC will not be liable for any loss, damage, cost or expense incurred or arising by reason of any person using or relying on the information in this publication.

CAUTION: RESEARCH ON UNREGISTERED AGRICULTURAL CHEMICAL USE

Any research with unregistered agricultural chemicals or of unregistered products reported in this document does not constitute a recommendation for that particular use by the authors or the authors' organisations.

All agricultural chemical applications must accord with the currently registered label for that particular agricultural chemical, crop, pest and region.

Copyright © All material published in this Fact Sheet is copyright protected and may not be reproduced in any form without written permission from the GRDC.

MORE INFORMATION

Dr Steve Marcroft

0409 978 941

steve@grainspathology.com.au

Dr Kurt Lindbeck

02 6938 1608

kurt.lindbeck@dpi.nsw.gov.au

Dr Ravjit Khangura

Ravjit.khangura@agric.wa.gov.au

0488 146 607

Mr Andrew Ware

Andrew.Ware@sa.gov.au

0427 884 272

PROJECT CODE: MGP0004